

Special Local Need Registration - 24(c)
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

Axiom[®] DF

Herbicide

EPA Reg. No. 3125-488

EPA SLN No. ID-010019

For use on established perennial grasses grown for seed.

ACTIVE INGREDIENTS:

Flufenacet54.4%
Metribuzin:13.6%

Restrictions

Do not use on clay knobs or poorly covered subsoils. Do not use on sand soils, or on any soils with less than 1% organic matter or on soils with pH higher than 7.5.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at time of pesticide application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Weeds controlled or suppressed

Annual Bluegrass
Bromus*
Italian (annual) ryegrass
Rattail Fescue
Volunteers of the established crop

* indicates suppression.

Special Precautions:

Do not apply this product through any type of irrigation equipment. Do not apply in tank-mixture with postemergence herbicides, such as Diuron. Application of oxyfluorfen or photosynthetic inhibiting postemergence herbicides within 4 weeks of AXIOM applications may cause crop damage.

Weed control may be reduced if excessive straw from the previous crop is present at AXIOM application or if insufficient rainfall/irrigation occurs after application.

Use only on fields which are at least one year old or have had at least one seed harvest. Use on perennial ryegrass after the second harvest may result in crop injury.

Recommended Applications

Crop	AXIOM DF Rate (ounces /acre)
Medium and fine textured soils greater than 1.5% organic matter	
Kentucky Bluegrass Tall Fescue Fine Fescue Perennial Ryegrass	9 to 13
Loamy sand and sandy loam soils greater than 1% organic matter	
Tall Fescue Fine Fescue Perennial Ryegrass	6 to 9

REMARKS : Apply specified dosage as a broadcast spray prior to weed seed emergence from the soil to no later than the 1 to 2 leaf stage of the volunteer grasses. Apply by ground equipment in 10 or more gallons of spray solution per acre. Applications made after mid-November may result in excessive crop injury and/or failure to control weeds. Use the higher rate of AXIOM under the following conditions: heavy surface vegetation residues or heavy weed pressure.

Allow at least 120 days between application and harvest.
Do not make more than one application per year.
Do not allow livestock to graze in treated areas.
Do not apply by aerial application.

IDAXSL01_027 11/09/01

Bayer Corporation
Crop Protection
Box 4913
Kansas City, MO 64120-0013
(800) 842-8020
<http://uscrop.bayer.com>

IMPORTANT

Before using this product, read and carefully observe the directions, cautionary statements and other information appearing on the product packaging label. This product is sold subject to the Conditions of Sale set forth on the container label.

